

LIGO-P070105-P-Z
Search for Gravitational Wave Bursts from Soft Gamma Repeaters

B. Abbott,¹⁶ R. Abbott,¹⁶ R. Adhikari,¹⁶ P. Ajith,² B. Allen,^{2, 54} G. Allen,³² R. Amin,²⁰ S. B. Anderson,¹⁶ W. G. Anderson,⁵⁴ M. A. Arain,⁴¹ M. Araya,¹⁶ H. Armandula,¹⁶ P. Armor,⁵⁴ Y. Aso,¹⁰ S. Aston,⁴⁰ P. Aufmuth,¹⁵ C. Aulbert,² S. Babak,¹ S. Ballmer,¹⁶ H. Bantilan,⁸ B. C. Barish,¹⁶ C. Barker,¹⁸ D. Barker,¹⁸ B. Barr,⁴² P. Barriga,⁵³ M. A. Barton,⁴² M. Bastarrika,⁴² K. Bayer,¹⁷ J. Betzwieser,¹⁶ P. T. Beyersdorf,²⁸ I. A. Bilenko,²³ G. Billingsley,¹⁶ R. Biswas,⁵⁴ E. Black,¹⁶ K. Blackburn,¹⁶ L. Blackburn,¹⁷ D. Blair,⁵³ B. Bland,¹⁸ T. P. Bodiya,¹⁷ L. Bogue,¹⁹ R. Bork,¹⁶ V. Boschi,¹⁶ S. Bose,⁵⁵ P. R. Brady,⁵⁴ V. B. Braginsky,²³ J. E. Brau,⁴⁷ M. Brinkmann,² A. Brooks,¹⁶ D. A. Brown,³³ G. Brunet,¹⁷ A. Bullington,³² A. Buonanno,⁴³ O. Burmeister,² R. L. Byer,³² L. Cadonati,⁴⁴ G. Cagnoli,⁴² J. B. Camp,²⁴ J. Cannizzo,²⁴ K. Cannon,¹⁶ J. Cao,¹⁷ L. Cardenas,¹⁶ T. Casebolt,³² G. Castaldi,⁵⁰ C. Cepeda,¹⁶ E. Chalkley,⁴² P. Charlton,⁹ S. Chatterji,¹⁶ S. Chelkowski,⁴⁰ Y. Chen,^{6, 1} N. Christensen,⁸ D. Clark,³² J. Clark,⁴² T. Cokelaer,⁷ R. Conte,⁴⁹ D. Cook,¹⁸ T. Corbitt,¹⁷ D. Coyne,¹⁶ J. D. E. Creighton,⁵⁴ A. Cumming,⁴² L. Cunningham,⁴² R. M. Cutler,⁴⁰ J. Dalrymple,³³ K. Danzmann,^{15, 2} G. Davies,⁷ D. DeBra,³² J. Degallaix,¹ M. Degree,³² V. Dergachev,⁴⁵ S. Desai,³⁴ R. DeSalvo,¹⁶ S. Dhurandhar,¹⁴ M. Díaz,³⁶ J. Dickson,⁴ A. Dietz,⁷ F. Donovan,¹⁷ K. L. Dooley,⁴¹ E. E. Doomes,³¹ R. W. P. Drever,⁵ I. Duke,¹⁷ J.-C. Dumas,⁵³ R. J. Dupuis,¹⁶ J. G. Dwyer,¹⁰ C. Echols,¹⁶ A. Effler,¹⁸ P. Ehrens,¹⁶ E. Espinoza,¹⁶ T. Etzel,¹⁶ T. Evans,¹⁹ S. Fairhurst,⁷ Y. Fan,⁵³ D. Fazi,¹⁶ H. Fehrmann,² M. M. Fejer,³² L. S. Finn,³⁴ K. Flasch,⁵⁴ N. Fotopoulos,⁵⁴ A. Freise,⁴⁰ R. Frey,⁴⁷ T. Fricke,^{16, 48} P. Fritschel,¹⁷ V. V. Frolov,¹⁹ M. Fyffe,¹⁹ J. Garofoli,¹⁸ I. Gholami,¹ J. A. Giaime,^{19, 20} S. Giampanis,⁴⁸ K. D. Giardino,¹⁹ K. Goda,¹⁷ E. Goetz,⁴⁵ L. Goggin,¹⁶ G. González,²⁰ S. Gossler,² R. Gouaty,²⁰ A. Grant,⁴² S. Gras,⁵³ C. Gray,¹⁸ M. Gray,⁴ R. J. S. Greenhalgh,²⁷ A. M. Gretarsson,¹¹ F. Grimaldi,¹⁷ R. Grosso,³⁶ H. Grote,² S. Grunewald,¹ M. Guenther,¹⁸ E. K. Gustafson,¹⁶ R. Gustafson,⁴⁵ B. Hage,¹⁵ J. M. Hallam,⁴⁰ D. Hammer,⁵⁴ C. Hanna,²⁰ J. Hanson,¹⁹ J. Harms,² G. Harry,¹⁷ E. Harstad,⁴⁷ K. Hayama,³⁶ T. Hayler,²⁷ J. Heefner,¹⁶ I. S. Heng,⁴² M. Hennessy,³² A. Heptonstall,⁴² M. Hewitson,² S. Hild,⁴⁰ E. Hirose,³³ D. Hoak,¹⁹ D. Hosken,³⁹ J. Hough,⁴² S. H. Huttner,⁴² D. Ingram,¹⁸ M. Ito,⁴⁷ A. Ivanov,¹⁶ B. Johnson,¹⁸ W. W. Johnson,²⁰ D. I. Jones,⁵¹ G. Jones,⁷ R. Jones,⁴² L. Ju,⁵³ P. Kalmus,¹⁰ V. Kalogera,²⁶ S. Kamat,¹⁰ J. Kanner,⁴³ D. Kasprzyk,⁴⁰ E. Katsavounidis,¹⁷ K. Kawabe,¹⁸ S. Kawamura,²⁵ F. Kawazoe,²⁵ W. Kells,¹⁶ D. G. Keppel,¹⁶ F. Ya. Khalili,²³ R. Khan,¹⁰ E. Khazanov,¹³ C. Kim,²⁶ P. King,¹⁶ J. S. Kissel,²⁰ S. Klimentenko,⁴¹ K. Kokeyama,²⁵ V. Kondrashov,¹⁶ R. K. Koppurapu,³⁴ D. Kozak,¹⁶ I. Kozhevnikov,¹³ B. Krishnan,¹ P. Kwee,¹⁵ P. K. Lam,⁴ M. Landry,¹⁸ M. M. Lang,³⁴ B. Lantz,³² A. Lazzarini,¹⁶ M. Lei,¹⁶ N. Leindegger,³² V. Leonhardt,²⁵ I. Leonor,⁴⁷ K. Libbrecht,¹⁶ H. Lin,⁴¹ P. Lindquist,¹⁶ N. A. Lockerbie,⁵² D. Lodhia,⁴⁰ M. Lormand,¹⁹ P. Lu,³² M. Lubinski,¹⁸ A. Lucianetti,⁴¹ H. Lück,^{15, 2} B. Machenschalk,² M. MacInnis,¹⁷ M. Mageswaran,¹⁶ K. Mailand,¹⁶ V. Mandic,⁴⁶ S. Márka,¹⁰ Z. Márka,¹⁰ A. Markosyan,³² J. Markowitz,¹⁷ E. Maros,¹⁶ I. Martin,⁴² R. M. Martin,⁴¹ J. N. Marx,¹⁶ K. Mason,¹⁷ F. Matichard,²⁰ L. Matone,¹⁰ R. Matzner,³⁵ N. Mavalvala,¹⁷ R. McCarthy,¹⁸ D. E. McClelland,⁴ S. C. McGuire,³¹ M. McHugh,²² G. McIntyre,¹⁶ G. McIvor,³⁵ D. McKechnan,⁷ K. McKenzie,⁴ T. Meier,¹⁵ A. Melissinos,⁴⁸ G. Mendell,¹⁸ R. A. Mercer,⁴¹ S. Meshkov,¹⁶ C. J. Messenger,² D. Meyers,¹⁶ J. Miller,^{42, 16} J. Minelli,³⁴ S. Mitra,¹⁴ V. P. Mitrofanov,²³ G. Mitselmakher,⁴¹ R. Mittleman,¹⁷ O. Miyakawa,¹⁶ B. Moe,⁵⁴ S. Mohanty,³⁶ G. Moreno,¹⁸ K. Mossavi,² C. MowLowry,⁴ G. Mueller,⁴¹ S. Mukherjee,³⁶ H. Mukhopadhyay,¹⁴ H. Müller-Ebhardt,² J. Munch,³⁹ P. Murray,⁴² E. Myers,¹⁸ J. Myers,¹⁸ T. Nash,¹⁶ J. Nelson,⁴² G. Newton,⁴² A. Nishizawa,²⁵ K. Numata,²⁴ J. O'Dell,²⁷ G. Ogin,¹⁶ B. O'Reilly,¹⁹ R. O'Shaughnessy,³⁴ D. J. Ottaway,¹⁷ R. S. Ottens,⁴¹ H. Overmier,¹⁹ B. J. Owen,³⁴ Y. Pan,⁴³ C. Pankow,⁴¹ M. A. Papa,^{1, 54} V. Parameshwaraiah,¹⁸ P. Patel,¹⁶ M. Pedraza,¹⁶ S. Penn,¹² A. Perreca,⁴⁰ T. Petrie,³⁴ I. M. Pinto,⁵⁰ M. Pitkin,⁴² H. J. Pletsch,² M. V. Plissi,⁴² F. Postiglione,⁴⁹ M. Principe,⁵⁰ R. Prix,² V. Quetschke,⁴¹ F. Raab,¹⁸ D. S. Rabeling,⁴ H. Radkins,¹⁸ N. Rainer,² M. Rakhmanov,³⁰ M. Ramsunder,³⁴ H. Rehbein,² S. Reid,⁴² D. H. Reitze,⁴¹ R. Riesen,¹⁹ K. Riles,⁴⁵ B. Rivera,¹⁸ N. A. Robertson,^{16, 42} C. Robinson,⁷ E. L. Robinson,⁴⁰ S. Roddy,¹⁹ A. Rodriguez,²⁰ A. M. Rogan,⁵⁵ J. Rollins,¹⁰ J. D. Romano,³⁶ J. Romie,¹⁹ R. Route,³² S. Rowan,⁴² A. Rüdiger,² L. Ruet,¹⁷ P. Russell,¹⁶ K. Ryan,¹⁸ S. Sakata,²⁵ M. Samidi,¹⁶ L. Sancho de la Jordana,³⁸ V. Sandberg,¹⁸ V. Sannibale,¹⁶ S. Saraf,²⁹ P. Sarin,¹⁷ B. S. Sathyaprakash,⁷ S. Sato,²⁵ P. R. Saulson,³³ R. Savage,¹⁸ P. Savov,⁶ S. W. Schediwy,⁵³ R. Schilling,² R. Schnabel,² R. Schofield,⁴⁷ B. F. Schutz,^{1, 7} P. Schwinberg,¹⁸ S. M. Scott,⁴ A. C. Searle,⁴ B. Sears,¹⁶ F. Seifert,² D. Sellers,¹⁹ A. S. Sengupta,¹⁶ P. Shawhan,⁴³ D. H. Shoemaker,¹⁷ A. Sibley,¹⁹ X. Siemens,⁵⁴ D. Sigg,¹⁸ S. Sinha,³² A. M. Sintes,^{38, 1} B. J. J. Slagmolen,⁴ J. Slutsky,²⁰ J. R. Smith,³³ M. R. Smith,¹⁶ N. D. Smith,¹⁷ K. Somiya,^{2, 1} B. Sorazu,⁴² L. C. Stein,¹⁷ A. Stochino,¹⁶ R. Stone,³⁶ K. A. Strain,⁴² D. M. Strom,⁴⁷ A. Stuver,¹⁹ T. Z. Summerscales,³ K.-X. Sun,³² M. Sung,²⁰ P. J. Sutton,⁷ H. Takahashi,¹ D. B. Tanner,⁴¹

R. Taylor,¹⁶ R. Taylor,⁴² J. Thacker,¹⁹ K. A. Thorne,³⁴ K. S. Thorne,⁶ A. Thüring,¹⁵ K. V. Tokmakov,⁴² C. Torres,¹⁹ C. Torrie,⁴² G. Traylor,¹⁹ M. Trias,³⁸ W. Tyler,¹⁶ D. Ugolini,³⁷ J. Ulmen,³² K. Urbanek,³² H. Vahlbruch,¹⁵ C. Van Den Broeck,⁷ M. van der Sluys,²⁶ S. Vass,¹⁶ R. Vaulin,⁵⁴ A. Vecchio,⁴⁰ J. Veitch,⁴⁰ P. Veitch,³⁹ A. Villar,¹⁶ C. Vorvick,¹⁸ S. P. Vyachanin,²³ S. J. Waldman,¹⁶ L. Wallace,¹⁶ H. Ward,⁴² R. Ward,¹⁶ M. Weinert,² A. Weinstein,¹⁶ R. Weiss,¹⁷ S. Wen,²⁰ K. Wette,⁴ J. T. Whelan,¹ S. E. Whitcomb,¹⁶ B. F. Whiting,⁴¹ C. Wilkinson,¹⁸ P. A. Willems,¹⁶ H. R. Williams,³⁴ L. Williams,⁴¹ B. Willke,^{15,2} I. Wilmot,²⁷ W. Winkler,² C. C. Wipf,¹⁷ A. G. Wiseman,⁵⁴ G. Woan,⁴² R. Wooley,¹⁹ J. Worden,¹⁸ W. Wu,⁴¹ I. Yakushin,¹⁹ H. Yamamoto,¹⁶ Z. Yan,⁵³ S. Yoshida,³⁰ M. Zanolin,¹¹ J. Zhang,⁴⁵ L. Zhang,¹⁶ C. Zhao,⁵³ N. Zotov,²¹ M. Zucker,¹⁷ and J. Zweizig¹⁶

(The LIGO Scientific Collaboration, <http://www.ligo.org>)

S. Barthelmy,²⁴ N. Gehrels,²⁴ K. C. Hurley,⁵⁶ and D. Palmer⁵⁷

¹Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-14476 Golm, Germany

²Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-30167 Hannover, Germany

³Andrews University, Berrien Springs, MI 49104 USA

⁴Australian National University, Canberra, 0200, Australia

⁵California Institute of Technology, Pasadena, CA 91125, USA

⁶Caltech-CaRT, Pasadena, CA 91125, USA

⁷Cardiff University, Cardiff, CF24 3AA, United Kingdom

⁸Carleton College, Northfield, MN 55057, USA

⁹Charles Sturt University, Wagga Wagga, NSW 2678, Australia

¹⁰Columbia University, New York, NY 10027, USA

¹¹Embry-Riddle Aeronautical University, Prescott, AZ 86301 USA

¹²Hobart and William Smith Colleges, Geneva, NY 14456, USA

¹³Institute of Applied Physics, Nizhny Novgorod, 603950, Russia

¹⁴Inter-University Centre for Astronomy and Astrophysics, Pune - 411007, India

¹⁵Leibniz Universität Hannover, D-30167 Hannover, Germany

¹⁶LIGO - California Institute of Technology, Pasadena, CA 91125, USA

¹⁷LIGO - Massachusetts Institute of Technology, Cambridge, MA 02139, USA

¹⁸LIGO Hanford Observatory, Richland, WA 99352, USA

¹⁹LIGO Livingston Observatory, Livingston, LA 70754, USA

²⁰Louisiana State University, Baton Rouge, LA 70803, USA

²¹Louisiana Tech University, Ruston, LA 71272, USA

²²Loyola University, New Orleans, LA 70118, USA

²³Moscow State University, Moscow, 119992, Russia

²⁴NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA

²⁵National Astronomical Observatory of Japan, Tokyo 181-8588, Japan

²⁶Northwestern University, Evanston, IL 60208, USA

²⁷Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX United Kingdom

²⁸San Jose State University, San Jose, CA 95192, USA

²⁹Sonoma State University, Rohnert Park, CA 94928, USA

³⁰Southeastern Louisiana University, Hammond, LA 70402, USA

³¹Southern University and A&M College, Baton Rouge, LA 70813, USA

³²Stanford University, Stanford, CA 94305, USA

³³Syracuse University, Syracuse, NY 13244, USA

³⁴The Pennsylvania State University, University Park, PA 16802, USA

³⁵The University of Texas at Austin, Austin, TX 78712, USA

³⁶The University of Texas at Brownsville and Texas Southmost College, Brownsville, TX 78520, USA

³⁷Trinity University, San Antonio, TX 78212, USA

³⁸Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain

³⁹University of Adelaide, Adelaide, SA 5005, Australia

⁴⁰University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁴¹University of Florida, Gainesville, FL 32611, USA

⁴²University of Glasgow, Glasgow, G12 8QQ, United Kingdom

⁴³University of Maryland, College Park, MD 20742 USA

⁴⁴University of Massachusetts, Amherst, MA 01003 USA

⁴⁵University of Michigan, Ann Arbor, MI 48109, USA

⁴⁶University of Minnesota, Minneapolis, MN 55455, USA

⁴⁷University of Oregon, Eugene, OR 97403, USA

⁴⁸University of Rochester, Rochester, NY 14627, USA

⁴⁹University of Salerno, 84084 Fisciano (Salerno), Italy

⁵⁰University of Sannio at Benevento, I-82100 Benevento, Italy

⁵¹University of Southampton, Southampton, SO17 1BJ, United Kingdom

⁵²University of Strathclyde, Glasgow, G1 1XQ, United Kingdom

⁵³University of Western Australia, Crawley, WA 6009, Australia

⁵⁴University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

⁵⁵Washington State University, Pullman, WA 99164, USA

⁵⁶University of California-Berkeley, Space Sciences Lab, 7 Gauss Way, Berkeley, CA 94720, USA

⁵⁷Los Alamos National Laboratory, Los Alamos, NM, 87545, USA

We present the results of a LIGO search for short-duration gravitational waves (GWs) associated with Soft Gamma Repeater (SGR) bursts. This is the first search sensitive to neutron star f -modes, usually considered the most efficient GW emitting modes. We find no evidence of GWs associated with any SGR burst in a sample consisting of the 27 Dec. 2004 giant flare from SGR 1806–20 and 190 lesser events from SGR 1806–20 and SGR 1900+14 which occurred during the first year of LIGO’s fifth science run. GW strain upper limits and model-dependent GW emission energy upper limits are estimated for individual bursts using a variety of simulated waveforms. The unprecedented sensitivity of the detectors allows us to set the most stringent limits on transient GW amplitudes published to date. We find upper limit estimates on the model-dependent isotropic GW emission energies (at a nominal distance of 10 kpc) between 3×10^{45} and 9×10^{52} erg depending on waveform type, detector antenna factors and noise characteristics at the time of the burst. These upper limits are within the theoretically predicted range of some SGR models.

Soft Gamma Repeaters (SGRs) sporadically emit brief (≈ 0.1 s) intense bursts of soft gamma-rays with peak luminosities commonly up to 10^{42} erg/s [1, 2]. Less common intermediate bursts with greater peak luminosities can last for seconds. Rare “giant flare” events, some 1000 times brighter than common bursts [3], have initial bright, short (≈ 0.2 s) pulses followed by tails lasting minutes and are among the most electromagnetically luminous events in the Universe [2]. Since the discovery of SGRs in 1979 three of the five confirmed SGRs have produced a giant flare each [4–7].

SGRs are promising sources of gravitational waves (GWs). According to the “magnetar” model SGRs are neutron stars with exceptionally strong magnetic fields $\sim 10^{15}$ G [8]. SGR bursts may result from the interaction of the star’s magnetic field with its solid crust, leading to crustal deformations and occasional catastrophic cracking [9, 10] with subsequent excitation of the star’s non-radial modes [11–13] and the emission of GWs [12–14]. Excitation of nonradial modes could also occur if SGRs are instead solid quark stars [14–16].

We present a search for short-duration GW signals ($\lesssim 0.3$ s) associated with SGR bursts using data collected by the Laser Interferometer Gravitational Wave Observatory (LIGO) [17]. LIGO consists of two co-located GW detectors at Hanford, WA with baselines of 4 km and 2 km and one 4 km detector at Livingston, LA. GW data from one or two of these detectors are used. When three detectors are operating, data from the most sensitive pair are chosen.

The SGR burst sample was provided by gamma-ray satellites of the interplanetary network [18], and includes the 27 Dec. 2004 giant flare from SGR 1806–20 and 214 confirmed bursts (152 from SGR 1806–20 and 62 from SGR 1900+14, one of which was a multi-episodic “storm” [19] occurring during the first year of LIGO’s fifth science run (S5) from 14 Nov. 2005 to 14 Nov. 2006.

Of the 214 bursts, 117 occurred with three LIGO detectors operating, 53 with two detectors operating, 20 with a single detector operating, and 24 with no detector operating. Including the giant flare, analysis was possible for a total of 191 listed SGR events.

To analyze a given SGR burst we divide the GW data into an on-source time region, in which GWs associated with the burst could be expected, and a background time region. In the background region we do not expect a GW associated with the SGR burst, but the noise is statistically similar to the on-source region. For isolated bursts the on-source region consists of 4 s of data centered on the SGR burst. GW emission is expected to occur almost simultaneously with the electromagnetic burst [13]; the 4 s on-source duration accounts for uncertainties in the geocentric electromagnetic peak time. There are three special cases: 1) for two SGR 1900+14 bursts which occurred within 4 s of each other a combined 7 s on-source region was chosen; 2) for the SGR 1900+14 storm a 40 s on-source region was used; 3) for an SGR 1806–20 event on 6 Aug. 2006 (hereafter 060806), two 4 s on-source regions were used, centered on the two distinct bright bursts comprising the event. Background regions consist of 1000 s of good data on either side of on-source regions. On-source and background segments are analyzed identically, including data quality cuts, resulting in lists of “analysis events.” Analysis events from the background regions are used to estimate the significance of the on-source analysis events; significant events, if any, are subject to environmental vetoes and consistency checks.

The analysis is performed by the *Flare pipeline* [20–22] and is based on the excess power detection statistic of [23]. Search parameters such as frequency bands and time windows are chosen to optimally detect the target signals. This is achieved by comparing detection efficiencies for simulated target signals injected into the background data and searched for with different search pa-

rameters [21]. The search targets neutron star fundamental mode ringdowns (RDs) predicted in [11–13, 24, 25] as well as unmodeled short-duration GW signals. Model predictions from [26] for ten realistic neutron star equations of state give f -mode RD frequencies in the range 1.5–3 kHz and damping times in the range 100–400 ms. We use a search band 1–3 kHz for RD searches (to include stiffer equations of state), and find a 250 ms time window to be optimal. The search for unmodeled signals uses time windows set by prompt SGR burst timescales (5–200 ms) and frequency bands set by the detector’s sensitivity; a 125 ms time window effectively covers this duration range, and we search in two bands: 100–200 Hz (probing the region in which the detectors are most sensitive) and 100–1000 Hz (for full spectral coverage below the ringdown search band).

In the absence of a detection, for each SGR burst we estimate loudest event upper limits [27] on the GW strain incident on the detector, h_{rSS} . Following [28] $h_{\text{rSS}}^2 = h_{\text{rSS}+}^2 + h_{\text{rSS}\times}^2$, where e.g. $h_{\text{rSS}+}^2 = \int_{-\infty}^{\infty} h_{+}^2 dt$ and $h_{+,\times}(t)$ are the two GW polarizations. The relationship between the GW polarizations and the detector response $h(t)$ to an impinging GW from an altitude and azimuth (θ, ϕ) and with polarization angle ψ is:

$$h(t) = F^{+}(\theta, \phi, \psi)h_{+}(t) + F^{\times}(\theta, \phi, \psi)h_{\times}(t) \quad (1)$$

where $F^{+}(\theta, \phi, \psi)$ and $F^{\times}(\theta, \phi, \psi)$ are the antenna functions for the source at (θ, ϕ) [29]. The upper limit is computed in a frequentist framework following the commonly used procedure of injecting simulated signals in the data and recovering them using the search pipeline (see for example [30, 31]). The upper limits are derived for RD signals and for unmodeled bursts. Correspondingly, RD and band-limited white noise burst (WNB) waveforms are injected with parameters chosen to probe the respective target signal space. For WNBs independent polarization components are generated with $h_{\text{rSS}+} = h_{\text{rSS}\times}$. For RD signals linearly and circularly polarized waves are considered. The polarization angle for each simulation was randomly chosen from a flat distribution between 0 and 2π .

The GW strain h_{rSS} upper limits can be recast as upper limits on the emitted GW energy, E_{GW} . Assuming isotropic emission, the GW energy associated with $h_{+}(t)$ and $h_{\times}(t)$ is [32]:

$$E_{\text{GW}} = 4\pi R^2 \frac{c^3}{16\pi G} \int_{-\infty}^{\infty} \left((\dot{h}_{+})^2 + (\dot{h}_{\times})^2 \right) dt. \quad (2)$$

We use this equation with a nominal source distance of $R = 10$ kpc (source locations and distances are discussed in [33, 34]) to compute the energies associated with the h_{rSS} upper limits for different signals.

Results — We find no evidence for gravitational waves associated with any of the SGR burst events in the sample. The significance of on-source analysis events is inferred by assigning rates at which background analysis

events of equal or greater loudness occur. We find the most significant on-source analysis event occurs at a rate of 1.35×10^{-3} Hz (1 per 741 s), which is consistent with the expectation for the 803 s of on-source data in the sample. We estimate 90% confidence strain and energy upper limits, $h_{\text{rSS}}^{90\%}$ and $E_{\text{GW}}^{90\%}$, using the loudest on-source analysis event for each SGR burst. Upper limits depend on detector sensitivity and antenna factors at the time of the burst, the loudest on-source analysis event, and the simulation waveform type used.

Table I lists upper limits for the SGR 1806–20 giant flare and for the brightest peak of the 060806 event from SGR 1806–20 [35] (complete results are given in [36]). Results from these two events are highlighted because they yield the smallest values of $\gamma = E_{\text{GW}}^{90\%}/E_{\text{EM}}$, a measure of the extent to which an energy upper limit probes the GW emission efficiency. At the time of the giant flare the LIGO Hanford 4 km detector was operating during a commissioning period (LIGO Astrowatch) and had noise amplitude higher than that of S5 by a factor of 3; the rms antenna factor, which is an indicator of the average sensitivity to a given source in the sky, for such event was $(F_{+}^2 + F_{\times}^2)^{1/2} = 0.3$. The isotropic electromagnetic energy (E_{EM}) for the event, assuming a distance of 10 kpc, was 1.6×10^{46} erg [6]. At the time of 060806 both the 4 km and 2 km Hanford detectors were observing, with rms antenna factor for that event of 0.5. E_{EM} for the brightest peak of 060806 was at least 2.9×10^{42} erg [35].

We estimate upper limits on GW strain and isotropic GW energy emitted using RDs with $\tau = 200$ ms at various frequencies, and WNBs lasting 11 and 100 ms and with 100–200 and 100–1000 Hz bands. We observe no more than 15% degradation in strain upper limits using RDs with τ in the range 100–300 ms, and no more than 20% degradation using WNBs with durations in the range 5–200 ms, as compared to the upper limits obtained for the nominal RDs and WNBs used for tuning the search. Superscripts in Table I give a systematic error and uncertainties at 90% confidence. The first and second superscripts account for systematic error and statistical uncertainty in amplitude and phase of the detector calibrations, estimated via Monte Carlo simulations, respectively. The third is a statistical uncertainty arising from using a finite number of injected simulations, estimated with the bootstrap method using 200 ensembles [37]. The systematic error and the quadrature sum of the statistical uncertainties are added to the final upper limit estimates.

Figure 1 shows $E_{\text{GW}}^{90\%}$ limits for the entire SGR burst sample. The lowest upper limit in the sample, $E_{\text{GW}}^{90\%} = 2.9 \times 10^{45}$ erg, is obtained for an SGR 1806–20 burst on 21 Jul. 2006, with a geocentric crossing time of 17:10:56.6 UTC. The lowest upper limit from the RD search is $E_{\text{GW}}^{90\%} = 2.4 \times 10^{48}$ erg for an SGR 1806–20 burst on 24 Aug. 2006 14:55:26 UTC.

Discussion — Two searches for GWs associated with




FIG. 1: $E_{\text{GW}}^{90\%}$ upper limits for the entire SGR burst sample for various circularly/linearly polarized RD (RDC/RDL) and white noise burst (WNB) signals. The limits shown in Table I, for the giant flare and the 060806 event, are indicated in the figure by circles and diamonds, respectively.

SGR events have been published previously; neither claimed detection. The AURIGA collaboration searched for GW bursts associated with the SGR 1806–20 giant flare in the band 850–950 Hz with damping time 100 ms, setting upper limits on the GW energy of $\sim 10^{49}$ erg [38]. The LIGO collaboration also published on the same giant flare, targeting times and frequencies of the quasi-periodic oscillations in the flare’s x-ray tail as well as other frequencies in the detector’s band, and setting upper limits on GW energy as low as 8×10^{46} erg for quasi-periodic signals lasting tens of seconds [39].

In addition to the 2004 giant flare, the search described here covers 190 lesser events which occurred during the LIGO S5 data run. Furthermore this search extends to the entire high sensitivity band of the detectors, which makes it the first search sensitive to neutron star f -modes, usually considered the most efficient GW emitting modes [11]. Our upper limits on E_{GW} overlap the range of E_{EM} 10^{44} – 10^{46} erg seen in SGR giant flares [3, 6]. Most of the WNB limits, and some of the RD limits, are below the 10^{49} erg maximum E_{GW} predicted in some theoretical models [13]. Our best upper limits on γ are within the theoretically predicted range implied in [13].

The Advanced LIGO detectors promise an improvement in h_{rss} by more than a factor of 10 over S5, corresponding to an improvement in energy sensitivity (and therefore γ) by more than a factor of 100. Thus within the next few years we expect to obtain GW energy upper limits for the f -mode search that fall in the E_{EM} range

of giant flares, and for the unmodeled search that fall in the E_{EM} range of intermediate bursts.

The authors are grateful to the Konus-Wind team and to S. Mereghetti for information used in the S5 burst list, and to G. Lichti and D. Smith for information on the giant flare event time. The authors gratefully acknowledge the support of the United States National Science Foundation for the construction and operation of the LIGO Laboratory and the Science and Technology Facilities Council of the United Kingdom, the Max-Planck-Society, and the State of Niedersachsen/Germany for support of the construction and operation of the GEO600 detector. The authors also gratefully acknowledge the support of the research by these agencies and by the Australian Research Council, the Council of Scientific and Industrial Research of India, the Istituto Nazionale di Fisica Nucleare of Italy, the Spanish Ministerio de Educación y Ciencia, the Conselleria d’Economia Hisenda i Innovació of the Govern de les Illes Balears, the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, The National Aeronautics and Space Administration, the Carnegie Trust, the Leverhulme Trust, the David and Lucile Packard Foundation, the Research Corporation, and the Alfred P. Sloan Foundation. K.C.H. is grateful for support under JPL Contracts 1282043 and 1268385, and NASA grants NAG5-11451 and NNG04GM50G. This paper is LIGO-P070105-05-Z.

-
- [1] S. Mereghetti, ArXiv e-prints **804** (2008), 0804.0250.
 - [2] P. M. Woods and C. Thompson, in *Compact Stellar X-Ray Sources*, edited by W. G. H. Lewin and M. van der Klis (Cambridge Univ. Press, Cambridge, 2004).
 - [3] D. M. Palmer et al., *Nature* **434**, 1107 (2005).
 - [4] E. P. Mazets et al., *Nature* **282**, 587 (1979).
 - [5] K. Hurley et al., *Nature* **397**, 41 (1999).
 - [6] K. Hurley et al., *Nature* **434**, 1098 (2005).
 - [7] S. Barthelmy et al., *GRB Coordinates Network* **8113** (2008).
 - [8] R. C. Duncan and C. Thompson, *Astrophys. J. Lett.* **392**, L9 (1992).
 - [9] C. Thompson and R. C. Duncan, *MNRAS* **275**, 255 (1995).
 - [10] S. J. Schwartz et al., *Astrophys. J. Lett.* **627**, L129 (2005).
 - [11] N. Andersson and K. D. Kokkotas, *MNRAS* **299**, 1059 (1998).
 - [12] J. A. de Freitas Pacheco, *Astron. and Astrophys.* **336**, 397 (1998).
 - [13] K. Ioka, *MNRAS* **327**, 639 (2001).
 - [14] J. E. Horvath, *Modern Phys. Lett. A* **20**, 2799 (2005).
 - [15] R. X. Xu, *Astrophys. J. Lett.* **596**, L59 (2003).
 - [16] B. J. Owen, *Phys. Rev. Lett.* **95**, 211101 (2005).
 - [17] LIGO Scientific Collaboration (2007), arXiv:0711.3041v1 [gr-qc].
 - [18] <http://ssl.berkeley.edu/ipn3>.

TABLE I: GW strain and energy upper limit estimates at 90% confidence ($h_{\text{rss}}^{90\%}$ and $E_{\text{GW}}^{90\%}$) for the SGR 1806–20 giant flare and the S5 SGR burst with the smallest limits on the ratio $\gamma = E_{\text{GW}}^{90\%}/E_{\text{EM}}$ for various circularly/linearly polarized RD (RDC/RDL) and white noise burst (WNB) waveforms. Uncertainties (given in superscripts for strain upper limits and explained in the text) are folded into the final upper limit estimates.

Waveform type	SGR 1806–20 Giant Flare			SGR 1806–20 060806		
	$h_{\text{rss}}^{90\%} [10^{-22} \text{ Hz}^{-\frac{1}{2}}]$	$E_{\text{GW}}^{90\%} [\text{erg}]$	γ	$h_{\text{rss}}^{90\%} [10^{-22} \text{ Hz}^{-\frac{1}{2}}]$	$E_{\text{GW}}^{90\%} [\text{erg}]$	γ
WNB 11ms 100-200 Hz	22 ^{+1.3 +5.6 +1.2} = 29	7.3×10^{47}	5×10^1	3.4 ^{+0.0 +0.4 +0.2} = 3.8	1.3×10^{46}	4×10^3
WNB 100ms 100-200 Hz	18 ^{+1.1 +4.6 +0.5} = 24	4.9×10^{47}	3×10^1	2.9 ^{+0.0 +0.3 +0.1} = 3.3	9.1×10^{45}	3×10^3
WNB 11ms 100-1000 Hz	50 ^{+3.0 +13 +1.3} = 66	5.4×10^{49}	3×10^3	7.5 ^{+0.0 +0.8 +0.3} = 8.3	8.3×10^{47}	3×10^5
WNB 100ms 100-1000 Hz	45 ^{+2.7 +12 +1.1} = 59	3.7×10^{49}	2×10^3	7.0 ^{+0.1 +0.7 +0.2} = 7.9	6.8×10^{47}	2×10^5
RDC 200ms 1090 Hz	59 ^{+3.6 +15 +1.7} = 78	2.6×10^{50}	2×10^4	10 ^{+0.2 +1.1 +0.4} = 12	5.8×10^{48}	2×10^6
RDC 200ms 1590 Hz	93 ^{+5.6 +24 +2.8} = 120	1.4×10^{51}	9×10^4	15 ^{+0.6 +1.5 +0.5} = 17	2.5×10^{49}	8×10^6
RDC 200ms 2090 Hz	120 ^{+7.4 +32 +3.5} = 160	4.2×10^{51}	3×10^5	20 ^{+1.6 +2.5 +0.6} = 24	8.9×10^{49}	3×10^7
RDC 200ms 2590 Hz	150 ^{+9.1 +39 +4.1} = 200	9.8×10^{51}	6×10^5	24 ^{+3.1 +3.0 +0.9} = 30	2.2×10^{50}	7×10^7
RDL 200ms 1090 Hz	170 ^{+10 +44 +36} = 240	2.6×10^{51}	2×10^5	33 ^{+1.0 +3.4 +3.5} = 38	6.7×10^{49}	2×10^7
RDL 200ms 1590 Hz	260 ^{+16 +68 +32} = 360	1.2×10^{52}	7×10^5	44 ^{+2.2 +4.6 +6.3} = 54	2.8×10^{50}	9×10^7
RDL 200ms 2090 Hz	390 ^{+23 +99 +46} = 520	4.4×10^{52}	3×10^6	64 ^{+7.0 +8.1 +9.1} = 83	1.1×10^{51}	4×10^8
RDL 200ms 2590 Hz	440 ^{+26 +110 +63} = 600	8.9×10^{52}	6×10^6	79 ^{+10 +10 +9.7} = 100	2.6×10^{51}	9×10^8

- [19] G. L. Israel et al., ArXiv e-prints **805** (2008), 0805.3919.
[20] P. Kalmus et al., *Class. Quant. Grav.* **24**, S659 (2007).
[21] P. Kalmus, PhD thesis, Columbia University (2008).
[22] P. Kalmus, mattapps Flare package, CVS tag ‘open4-v3’ (2008), http://www.lsc-group.phys.uwm.edu/cgi-bin/cvs/viewcvs.cgi/?cvsroot=1_scssoft.
[23] W. G. Anderson, P. R. Brady, J. D. E. Creighton, and É. É. Flanagan, *Phys. Rev. D* **63**, 042003 (2001).
[24] B. L. Schumaker and K. S. Thorne, *MNRAS* **203**, 457 (1983).
[25] N. Andersson, *Class. Quant. Grav.* **20**, 105 (2003).
[26] O. Benhar, V. Ferrari, and L. Gualtieri, *Phys. Rev. D* **70**, 124015 (2004).
[27] P. R. Brady, J. D. E. Creighton, and A. G. Wiseman, *Class. Quant. Grav.* **21** (2004).
[28] B. Abbott et al. (LIGO Scientific Collaboration), *Phys. Rev. D* **72**, 062001 (2005).
[29] K. S. Thorne, in *300 Years of Gravitation*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1987), p. 417.
[30] B. Abbott et al. (LIGO Scientific Collaboration), *Phys. Rev. D* **76**, 082001 (2007).
[31] B. Abbott et al. (LIGO Scientific Collaboration), *Phys. Rev. D* **77**, 062004 (2008).
[32] S. Shapiro and S. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars* (Wiley, New York, 1983).
[33] S. Corbel and S. S. Eikenberry, *Astron. Astrophys.* **419**, 191 (2004).
[34] D. L. Kaplan, S. R. Kulkarni, D. A. Frail, and M. H. van Kerkwijk, *Astrophys. J.* **566**, 378 (2002).
[35] S. Golenetskii et al., *GRB Coord. Network* **5426**, (2006).
[36] See EPAPS Document No. XX for complete table. For more information on EPAPS, see <http://www.aip.org/pubservs/epaps.html>.
[37] B. Efron, *Ann. Statist.* **7**, 1 (1979).
[38] L. Baggio et al., *Phys. Rev. Lett.* **95**, 081103 (2005); *Phys. Rev. Lett.* **95**, 139903(E) (2005).
[39] B. Abbott et al. (LIGO Scientific Collaboration), *Phys. Rev. D* **76**, 062003 (2007).