


First Results from LIGO

Keith Riles
University of Michigan



Physics and Astronomy
Colloquium

University of Rochester

January 19, 2005



Outline



- ❑ **Nature & Generation of Gravitational Waves**
- ❑ **Detecting Gravitational Waves with the LIGO Detector**
- ❑ **Data runs and Early Results**
- ❑ **Looking Ahead – Advanced LIGO**

Nature of Gravitational Waves

- Gravitational Waves = “Ripples in space-time”
- Perturbation propagation similar to light (obeys same wave equation!)
 - ◆ Propagation speed = c
 - ◆ Two transverse polarizations - quadrupolar: $+$ and \times

Example:

Ring of test masses
responding to wave
propagating along z



- Amplitude parameterized by (tiny)
dimensionless strain h : $\Delta L \sim h(t) \times L$


Why look for Gravitational Radiation?

- Because it's there! (presumably)

- Test General Relativity:
 - ◆ Quadrupolar radiation? Travels at speed of light?
 - ◆ Unique probe of strong-field gravity

- Gain different view of Universe:
 - ◆ Sources cannot be obscured by dust
 - ◆ Detectable sources some of the most interesting, least understood in the Universe
 - ◆ Opens up entirely new non-electromagnetic spectrum

What will the sky look like?



Generation of Gravitational Waves


- Radiation generated by quadrupolar mass movements:

$$h_{\mu\nu} = \frac{2G}{rc^4} \frac{d^2}{dt^2} (I_{\mu\nu})$$

(with $I_{\mu\nu}$ = quadrupole tensor, r = source distance)

- Example: Pair of $1.4 M_{\text{solar}}$ neutron stars in circular orbit of radius 20 km (imminent coalescence) at orbital frequency 400 Hz gives 800 Hz radiation of amplitude:

$$h \approx \frac{10^{-21}}{(r/15\text{Mpc})}$$



Generation of Gravitational Waves

Major expected sources in 10-1000 Hz “terrestrial” band:

- ❑ Coalescences of binary compact star systems
(NS-NS, NS-BH, BH-BH)
- ❑ Supernovae
(requires asymmetry in explosion)
- ❑ Spinning neutron stars, e.g., pulsars
(requires axial asymmetry or wobbling spin axis)

Also expected (but probably exceedingly weak):

- ❑ Stochastic background – Big Bang remnant


Or from Cosmic Strings? → Talk to Adrian!

Generation of Gravitational Waves

- ❑ Strong indirect evidence for GW generation:

Taylor-Hulse Pulsar System (PSR1913+16)

- ♦ Two neutron stars (one=pulsar) in elliptical 8-hour orbit
- ♦ Measured periastron advance quadratic in time in agreement with absolute GR prediction
→ **Orbital decay due to energy loss**




Generation of Gravitational Waves

Can we detect this radiation directly?


NO - freq too low

Must wait ~ 300 My for characteristic “chirp”:




Generation of Gravitational Waves


Audio



Last nine seconds of inspiral



First Res



Generation of Gravitational Waves

Coalescence rate estimates based on two methods:

- Use known NS/NS binaries in our galaxy (three!)
- *A priori* calculation from stellar and binary system evolution

→ Large uncertainties!

For initial LIGO design “seeing distance” (~20 Mpc):

Expect 1/(70 y) to 1/(4 y)

→ Will need Advanced LIGO to ensure detection

Generation of Gravitational Waves

Most promising periodic source: Rotating Neutron Stars (e.g., pulsar)

But axisymmetric object rotating about symmetry axis
Generates NO radiation

Need an asymmetry or perturbation:


- Equatorial ellipticity (e.g., – mm-high “mountain”):

$$h \propto \epsilon_{\text{equat}}$$

- Poloidal ellipticity (natural) + wobble angle (precessing star):

$$h \propto \epsilon_{\text{pol}} \times \Theta_{\text{wobble}}$$

(precession due to different L and Ω axes)



Periodic Sources of GW

Serious technical difficulty: Doppler frequency shifts

- ◆ Frequency modulation from earth's rotation ($v/c \sim 10^{-6}$)
- ◆ Frequency modulation from earth's orbital motion ($v/c \sim 10^{-4}$)

Additional, related complications:

- ◆ Daily amplitude modulation of antenna pattern
- ◆ Spin-down of source
- ◆ Orbital motion of sources in binary systems

Modulations / drifts complicate analysis enormously:

- ◆ Simple Fourier transform inadequate
- ◆ Every sky direction requires different demodulation
→ All-sky survey at full sensitivity = **Formidable challenge**

Periodic Sources of GW

But two substantial benefits from modulations:

- ◆ Reality of signal confirmed by need for corrections
- ◆ Corrections give precise direction of source

□ Difficult to detect spinning neutron stars!

□ But search is nonetheless intriguing:

- ◆ Unknown number of electromagnetically quiet, undiscovered neutron stars in our galactic neighborhood
- ◆ Realistic values for ϵ unknown
- ◆ A nearby source could be buried in the data, waiting for just the right algorithm to tease it into view

Much effort underway → Expect results in April


Outline

- ❑ **Nature & Generation of Gravitational Waves**
- ❑ **Detecting Gravitational Waves with the LIGO Detector**
- ❑ **Data runs and Early Results**
- ❑ **Preparing for Advanced LIGO**

Gravitational Wave Detection

□ Suspended Interferometers (IFO's)

- ◆ Suspended mirrors in “free-fall”
- ◆ Broad-band response
(~50 Hz to few kHz)
- ◆ Waveform information
(e.g., chirp reconstruction)
- ◆ Michelson IFO is
“natural” GW detector



Gravitational Wave Detection

Major Interferometers coming on line world-wide


LIGO (NSF-\$300M) Livingston, Louisiana & Hanford, Washington	2 x 4000-m 1 x 2000-m	Advanced Commissioning & Data Taking
VIRGO Near Pisa, Italy	1 x 3000-m	Early Commissioning
GEO Near Hannover, Germany	1 x 600-m	Advanced Commissioning & Data Taking
TAMA Tokyo, Japan	1 x 300-m	Advanced Commissioning & Data Taking

LIGO Interferometer Optical Scheme

Michelson interferometer

With Fabry-Perot arm cavities

• Recycling mirror matches losses, enhances effective power by ~ 50x



“Locking” the Inteferometer

Sensing gravitational waves requires sustained resonance in the Fabry-Perot arms and in the recycling cavity

- Need to maintain half-integer # of laser wavelengths between mirrors
- Feedback control servo uses error signals from imposed RF sidebands
- Four primary coupled degrees of freedom to control
- Highly non-linear system with 5-6 orders of magnitude in light intensity

Also need to control mirror rotation (“pitch” & “yaw”)

- Ten more DOF’s (but less coupled)

And need to stabilize laser (intensity & frequency), keep the beam pointed, damp out seismic noise, correct for tides, etc.,...


LIGO Observatories

Hanford




Observation of nearly simultaneous signals 3000 km apart rules out terrestrial artifacts

Livingston



LIGO Detector Facilities



- Stainless-steel tubes
(1.24 m diameter, $\sim 10^{-8}$ torr)
- Gate valves for optics isolation
- Protected by concrete enclosure

Vacuum System




LIGO Detector Facilities

LASER

- ❑ Infrared (1064 nm, 10-W) Nd-YAG laser from Lightwave (now commercial product!)
- ❑ Elaborate intensity & frequency stabilization system, including feedback from main interferometer

Optics


- ❑ Fused silica (high-Q, low-absorption, 1 nm surface rms, 25-cm diameter)
- ❑ Suspended by single steel wire
- ❑ Actuation of alignment / position via magnets & coils



LIGO Detector Facilities

Seismic Isolation

- ❑ Multi-stage (mass & springs) optical table support gives 10^6 suppression
- ❑ Pendulum suspension gives additional $1 / f^2$ suppression above ~ 1 Hz




What Limits the Sensitivity of the Interferometers?

- Seismic noise & vibration limit at low frequencies
- Atomic vibrations (Thermal Noise) inside components limit at mid frequencies
- Quantum nature of light (Shot Noise) limits at high frequencies
- Myriad details of the lasers, electronics, etc., can make problems above these levels

Best design sensitivity:

$\sim 3 \times 10^{-23} \text{ Hz}^{-1/2} @ 150 \text{ Hz}$



Some interesting problems at Hanford...

Brush fire sweeps over site
– June 2000




Charred landscape,
but no IFO damage!



Tacoma earthquake –
Feb 2001

- Misaligned optics
- Actuation magnets dislodged
- Commissioning delay



Human error too!

And a new problem to worry about...



**Mt. St. Helens
has awoken!**

**Micro-quakes in
late September
interfered with
commissioning**



**Eruption in
early October
helped –
relieved
pressure!**

Livingston Problem -- Logging



Livingston Observatory
located in pine forest popular
with pulp wood cutters

Spiky noise (e.g. falling trees) in
1-3 Hz band creates dynamic
range problem for arm cavity
control


→ **40% livetime**

Solution: Retrofit with active feed-forward isolation system
(using technology developed for Advanced LIGO)

→ Work started January 2004



→ Commissioning nearly complete – **Looks promising!**

LIGO Organization & Support



LIGO Scientific Collaboration

The LIGO Logo's



K. Riles - First Results from LIGO -11/30/04

GEO600

Work closely with the GEO600 Experiment (Germany / UK / Spain)

- **Arrange coincidence data runs when commissioning schedules permit**
- **GEO members are full members of the LIGO Scientific Collaboration**
- **Data exchange and strong collaboration in analysis now routine**
- **Major partners in proposed Advanced LIGO upgrade**



**600-meter Michelson Interferometer
just outside Hannover, Germany**

Outline

- ❑ **Nature & Generation of Gravitational Waves**
- ❑ **Detecting Gravitational Waves with the LIGO Detector**
- ❑ **Data runs and Early Results**
- ❑ **Looking Ahead – Advanced LIGO**

Data Runs

Have carried out a series of Engineering Runs (E1--E11) and Science Runs (S1--S3) interspersed with commissioning

S1 run:

17 days (August / September 2002)

Four detectors operating: LIGO (L1, H1, H2) and GEO600

H1 (235 hours) H2(298 hours) L1(170 hours)

Triple-LIGO-coincidence (96 hours)

Four S1 astrophysical searches published (Physical Review D):

» **Inspiring neutron stars -- PRD 69 (2004) 122001**

» **Bursts -- PRD 69 (2004) 102001**

» **Known pulsar (J1939+2134) – PRD 69 (2004) 082004**

» **Stochastic background -- PRD 69 (2004) 122004**

Data Runs

S2 run:

59 days (February—April 2003)

Four interferometers operating: LIGO (L1, H1, H2) and TAMA300 plus Allegro bar detector at LSU

H1 (1044 hours) H2 (822 hours) L1 (536 hours)

Triple-LIGO-coincidence (318 hours)

Many S2 searches underway – some prelim./final results for today:

» Inspiring neutron stars

» Coincidence with gamma ray burst GRB030329

» 28 known pulsars

» Stochastic background

S3 run:


70 days (October 2003 – January 2004) – Analysis underway...

S2 Sensitivities

Strain Sensivities for the LIGO Interferometers for S2

14 February 2003 - 14 April 2003

LIGO-G030379-00-E



Livingston (L1)
Interferometer
most sensitive
in "sweet spot"

Inspiring Neutron Stars – S2 Results

S2 sensitivity permitted seeing the Andromeda Galaxy with L1 whenever live, with H1 seeing it at times (when noise low and antenna pattern favorable)

Analysis based on matched filtering in Fourier domain (hundreds of templates in bank for $M_{\odot} < M_1, M_2 < 3 M_{\odot}$)

Inspiral triggers parameterized by signal-to-noise ratio and frequency-domain χ^2


Vetoed on L1 triggers coincident with auxiliary channel artifacts

“Playground” (10%) data used to tune thresholds, vetoed for remaining 90%


Hanford-Livingston coincidence required

Inspiring Neutron Stars

Background with simulated signals



Observed events



No evidence for excess events

→ Set limit based on “Loudest event statistic”



→ Obtain preliminary rate:

$R_{90\%} < 50$ inspirals per year per “milky-way-equivalent-galaxy”

K. Riles - First Results from LIGO -11/30/04

Search for “Generic” Bursts (S1 results)


(look for coincident pulses in time-freq plane)



Background rates measured from non-zero
time shifts between interferometers


Feldman-Cousins 90% CL upper limit: **< 1.6 events/day**

90% CL rate limit vs. strength plots for two burst models




Burst model:

1ms, 2.5 ms Gaussian impulses



Burst model:

Sine-Gaussians with varying central frequencies



- ❑ Determined **detection efficiency** via **signal injections**
- ❑ Assumed a **population** of such sources **uniformly** distributed on a concentric sphere

Gamma Ray Burst 030329 – S2 Results

GRB030329 was a powerful burst that occurred during the S2 run


Identified in gammas, x-rays, and optical

Spectroscopy strongly suggests Supernova origin:

Distance (800 Mpc!) made it unlikely to be detectable by LIGO, but event provides interesting “practice run” for GRB detection (L1 off at time ☹)

Supernova Spectrum Emergence

GRB 030329 is now also SN2003dh




T. Matheson (CfA), GCN 2120

Gamma Ray Burst 030329

Searched for excess cross-correlation between Hanford Interferometers


Examined background noise and set false alarm probability for 3-minute interval around GRB to be ~10%

Estimated efficiencies from generic (sine-Gaussian) signal injections for varying central frequencies & Q's




Simulation

Integration Length




Time 40

Gamma Ray Burst 030329




Expected numbers of events (using two different background estimates) and **observed** numbers of events vs “event strength”



No candidates above (or even near) threshold
 → Set upper limits:



Known Pulsars – S2 Results



Crab pulsar



- **Detectable amplitudes** with a 1% false alarm rate and 10% false dismissal rate by the IFOs during S2 (colored curves) and at design sensitivities (black curves)
- **Upper limits** on $\langle h_o \rangle$ from spin-down measurements of known radio pulsars (filled circles)

Searched for 28 known isolated pulsars for which precise timing information is available from radio astronomers

Known Pulsars

Search based on coherent time-domain heterodyne, accounting for Doppler shifts due to Earth's spin and orbital motion; and accounting for antenna pattern amplitude modulations

Can reconstruct amplitude, phase, polarization and orientation of strong source

Parameter fitting for hardware-injected fake pulsar during S2:

Posterior probability densities for PSR signal \mathcal{Z}





Flat priors for $\cos\psi$, ψ , ϕ_0 , h_0 ($h_0 > 0$); Jeffreys' prior for σ ($p(\sigma) \propto 1/\sigma$)

solid red line - L1

solid blue line - H2

solid green line - H1

dashed black line - Joint



Known Pulsars

No signals detected

Obtained upper limits on source strengths:

- Amplitudes h_0
- Pulsar ellipticities ε


based on inferred Bayesian posterior probability density functions (pdf's) (flat prior for h_0)

Best 95% CL upper limit on h_0 :

1.7×10^{-24} (J1910-5959D)

Sample pdf for the Crab pulsar (B0531+21)

→ 95% CL upper limit on h_0 : 4.1×10^{-22}



Stochastic Background – S2 Results

- **Sources:** early universe, many weak unresolved sources emitting gravitational waves independently
→ Random radiation described by its spectrum (isotropic, unpolarized, stationary and Gaussian)
- **Analysis goals:** constrain contribution of stochastic radiation's energy ρ_{GW} to the total energy required to close the universe ρ_{critical} :

$$\int_0^{\infty} (1/f) \Omega_{\text{GW}}(f) df = \frac{\rho_{\text{GW}}}{\rho_{\text{critical}}}$$

- Use optimally filtered **cross-correlation** of detector pairs: L1-H1, L1-H2 and H1-H2
→ Report L1-H1 results today

Stochastic Background

Detector separation and orientation reduce correlations at high frequencies

($\lambda_{\text{GW}} \geq 2 \times \text{BaseLine}$)


H1-H2 most sensitive
(but instruments correlated!)

L1-H1(H2) most sensitive < 50 hz

Known inter-site correlated lines removed in analysis

Assume simple model: $\Omega(f) = \Omega_0$

Cumulative measure of Ω_0 during the S2 run



Preliminary 90% CL limit:

$$\Omega_0 (h_{100})^2 < 0.017$$


Outline

- ❑ **Nature & Generation of Gravitational Waves**
- ❑ **Detecting Gravitational Waves with the LIGO Detector**
- ❑ **Data runs and Early Results**
- ❑ **Looking Ahead – Advanced LIGO**

Looking Ahead

Best Strain Sensitivities for the LIGO Interferometers

Comparisons among S1, S2, S3 LIGO-G030548-02-E



Looking Ahead

Resume data runs in February 2005:

- **Verify success of Livingston seismic retrofit**
- **Verify success of sensitivity improvements**

First true “Search Run” in late 2005


Plan before shutdown for Advanced LIGO upgrade:

≥ 1 year of running at Initial LIGO design sensitivity

Looking Ahead

The three LIGO and the GEO interferometers are part of a forming **Global Network**.

Multiple signal detections will increase detection confidence and provide better precision on source locations and wave polarizations




Looking Further Ahead

Despite their immense technical challenges, the initial LIGO IFO's were designed conservatively, based on “tabletop” prototypes, but with expected sensitivity gain of ~ 1000 .

Given the expected low rate of detectable GW events, it was always planned that in engineering, building and commissioning initial LIGO, one would learn how reliably to build Advanced LIGO with another factor of ~ 10 improved sensitivity.

Because LIGO measures GW amplitude, an increase in sensitivity by 10 gives an increase in sampling volume, i.e, rate by ~ 1000




Advanced LIGO

Sampling of source strengths vis a vis Initial LIGO and Advanced LIGO

Lower h_{rms} and wider bandwidth both important

“Signal recycling” offers potential for tuning shape of noise curve to improve sensitivity in target band (e.g., known pulsar cluster)



Advanced LIGO

Increased laser power:

10 W → 180 W

Improved shot noise (high freq)

Potential new test mass material:

Fused silica → Sapphire


Lower internal thermal noise in bandwidth

Increased test mass:

10 kg → 40 kg

Compensates increased radiation pressure noise

Sapphire Optics



Date: 10/25/2001
Time: 13:59:18
Wavelength: 1.064 um
Pupil: 100.0 %
PV: 81.6271 nm
RMS: 13.2016 nm

X Center: 172.00
Y Center: 145.00
Radius: 163.00 pix
Terms: None
Filters: None
Masks:



Advanced LIGO

Detector Improvements:

New suspensions:

Single → Quadruple pendulum

**Lower suspensions thermal noise
in bandwidth**



Improved seismic isolation:

Passive → Active

Lowers seismic “wall” to ~10 Hz

Conclusions

LIGO commissioning is well underway

- Good progress toward design sensitivity
- GEO, other instruments advancing as well

Science Running is beginning

- Initial results from our first two data runs

Our Plan:

- Continue commissioning and data runs with GEO & others
- Collect \geq one year of data at design sensitivity before starting upgrade
- Advanced interferometer with dramatically improved sensitivity – 2009+
(NSF MRE proposal recently approved by National Science Board)

**We should be detecting gravitational waves
routinely within the next 10 years!**