GW150914: FACTSHEET

BACKGROUND IMAGES: TIME-FREQUENCY TRACE (TOP) AND TIME-SERIES (BOTTOM) IN THE TWO LIGO DETECTORS; SIMULATION OF BLACK HOLE HORIZONS (MIDDLE-TOP), BEST FIT WAVEFORM (MIDDLE-BOTTOM)

first direct detection of gravitational waves (GW) and first direct observation of a black hole binary

observed by	LIGO L1, H1
source type	black hole (BH) binary
date	14 Sept 2015
time	09:50:45 UTC
likely distance	0.75 to 1.9 Gly 230 to 570 Mpc
redshift	0.054 to 0.136
signal-to-noise ratio	24
false alarm prob.	< 1 in 5 million
false alarm rate	< 1 in 200,000 yr

Source Masses	M⊙
total mass	60 to 70
primary BH	32 to 41
secondary BH	25 to 33
remnant BH	58 to 67
mass ratio	0.6 to 1
primary BH spin	< 0.7
secondary BH spin	< 0.9
remnant BH spin	0.57 to 0.72
signal arrival time delay	arrived in L1 7 ms before H1
likely sky position	Southern Hemisphere
likely orientation	face-on/off
resolved to	~600 sq. deg.

duration from 30 Hz	~ 200 ms
# cycles from 30 Hz	~10
peak GW strain	1×10^{-21}
peak displacement of interferometers arms	±0.002 fm
frequency/wavelength at peak GW strain	150 Hz, 2000 km
peak speed of BHs	~ 0.6 c
peak GW luminosity	3.6×10^{56} erg s^{-1}
radiated GW energy	2.5-3.5 M⊙
remnant ringdown freq.	~ 250 Hz
remnant damping time	~ 4 ms
remnant size, area	180 km, 3.5×10^5 km^2
consistent with general relativity?	passes all tests performed
graviton mass bound	< 1.2×10^{-22} eV
coalescence rate of binary black holes	2 to 400 Gpc^{-3} yr^{-1}
online trigger latency	~ 3 min
# offline analysis pipelines	5
CPU hours consumed	~ 50 million (=20,000 PCs run for 100 days)
papers on Feb 11, 2016	13
# researchers	~1000, 80 institutions in 15 countries

Detector noise introduces errors in measurement. Parameter ranges correspond to 90% credible bounds.
Acronyms: L1=LIGO Livingston, H1=LIGO Hanford; Gly=giga lightyear=9.46×10^{21} km; Mpc=mega parsec=3.2 million lightyear, Gpc=10^3 Mpc, fm=femtometer=10^{-15} m, M⊙=1 solar mass=2×10^{30} kg